

SLOW AREA-PRESERVING DIFFEOMORPHISMS OF THE TORUS

ALEXANDER BORICHEV

ABSTRACT. We construct area-preserving real analytic diffeomorphisms of the torus with unbounded growth sequences of arbitrarily slow growth.

Given a smooth compact manifold M , consider the group $\text{Diff}(M)$ of diffeomorphisms of M . For every $f \in \text{Diff}(M)$ we define the growth sequence of f :

$$\Gamma_n(f) = \max\left(\max_{x \in M} \|d_x f^n\|, \max_{x \in M} \|d_x f^{-n}\|\right), \quad n \in \mathbb{N},$$

where f^n is the n -th iteration of f , f^{-n} is the n -th iteration of f^{-1} , and $\|d_x f\|$ is the operator norm of the differential of f at the point $x \in M$. Conjugations of f in the group $\text{Diff}(M)$ generate equivalent growth sequences:

$$c(g)\Gamma_n(g^{-1}fg) \leq \Gamma_n(f) \leq C(g)\Gamma_n(g^{-1}fg), \quad g \in \text{Diff}(M), n \in \mathbb{N}.$$

The asymptotics of the growth sequence is a basic dynamic invariant (see [3]). D'Ambra and Gromov [1, 7.10.C] proposed to study the behavior of growth sequences for various classes of diffeomorphisms. In particular, it is interesting to find examples of unbounded growth sequences of slow growth (see also the references in [1, 7.10.C]). We call the diffeomorphisms generating such growth sequences the slow diffeomorphisms.

Recently, Polterovich and Sodin [5] obtained several results on the growth sequences of smooth order-preserving diffeomorphisms of the interval $[0, 1]$. In particular, they proved [5, Theorem 1.7] that for any sequence $\{a_n\}$ of positive numbers tending to infinity, there exists a C^∞ -smooth diffeomorphism f , $f \neq \text{Id}$, such that

$$\liminf_{n \rightarrow \infty} \frac{\Gamma_n(f)}{a_n} \leq 1.$$

Key words and phrases. Area-preserving diffeomorphisms, growth sequences.
 2000 *Mathematical Subject Classification* 37C05, 26A18, 28D05, 58D05.
E-mail: borichev@math.u-bordeaux.fr.

On the other hand, a simple argument (see [5]) shows that for any diffeomorphism $f \neq \text{Id}$,

$$\sum_{n \geq 1} \frac{1}{\Gamma_n(f)} < \infty.$$

Furthermore, Polterovich proved [4, Theorem 1.3] that for every $0 < \beta < 1$, there exists an area-preserving real analytic diffeomorphism f of the torus such that

$$\begin{aligned} \Gamma_n(f) &\leq Cn^\beta \log n, \\ \limsup_{n \rightarrow \infty} \frac{\Gamma_n(f)}{n^\beta} &> 0. \end{aligned}$$

In this note, we improve somewhat the result by Polterovich by producing area-preserving real analytic diffeomorphisms of the torus with arbitrarily slowly growing unbounded growth sequences.

Theorem. *Let φ be a positive increasing (unbounded) function on $\mathbb{R}_+$ such that $\varphi(x) = o(x)$, $x \rightarrow \infty$. There exists an area-preserving real analytic diffeomorphism f of the torus such that*

$$\left. \begin{aligned} \Gamma_n(f) &\leq \varphi(n), \\ \limsup_{n \rightarrow \infty} \frac{\Gamma_n(f)}{\varphi(n)} &> 0. \end{aligned} \right\} \quad (1)$$

For some related questions on the asymptotics of diffeomorphisms with fixed points see [4]. Other recent results on the behavior of the growth sequences are in [2], [5].

Proof of the Theorem. We represent the torus as the product $[0, 1) \times [0, 1)$, and define, as in [4],

$$f(x, y) = (\{x + \alpha\}, \{y + cF(x)\}), \quad x, y \in [0, 1),$$

for $\alpha \in \mathbb{R}$, $c \in (0, +\infty)$, and a real analytic 1-periodic function $F : \mathbb{R} \rightarrow \mathbb{R}$; here $\{\cdot\}$ stands for the fractional part. Then

$$d_x f = \begin{pmatrix} 1 & 0 \\ cF'(x) & 1 \end{pmatrix},$$

and f is an area-preserving real analytic diffeomorphism of the torus.

Define the Weyl sums

$$W(N, x, \alpha) = \sum_{n=0}^{N-1} F'(x + n\alpha). \quad (2)$$

We have

$$d_x f^N = \begin{pmatrix} 1 & 0 \\ cW(N, x, \alpha) & 1 \end{pmatrix}, \quad N \geq 1,$$

and

$$d_x f^{-N} = \begin{pmatrix} 1 & 0 \\ -cW(N, x - N\alpha, \alpha) & 1 \end{pmatrix}, \quad N \geq 1.$$

Therefore, for (1) to hold it suffices that F and α satisfy the following condition:

$$0 < \limsup_{N \rightarrow \infty} \max_{0 \leq x < 1} \frac{W(N, x, \alpha)}{\varphi(N)} < \infty. \quad (3)$$

Up to now our proof repeated that of Polterovich in [4]. The main difference of our argument is in the way of estimating the Weyl sums (2).

We are going to choose a sequence $\{q_k\}_{k \geq 1}$, $q_1 = 1$,

$$\frac{q_{k+1}}{100q_k} \in \mathbb{N}, \quad k \geq 1, \quad (4)$$

and a sequence $\{r_k\}_{k \geq 1}$,

$$0 < r_k < \exp(-q_k), \quad k \geq 1, \quad (5)$$

and define

$$F(x) = \sum_{k \geq 1} \frac{r_k}{2\pi q_k} \sin[2\pi q_k x], \quad x \in \mathbb{R}.$$

Then F is real analytic and 1-periodic,

$$F'(x) = \sum_{k \geq 1} r_k \cos[2\pi q_k x], \quad x \in \mathbb{R}.$$

For $\alpha \in \mathbb{R}$ denote

$$\begin{aligned} \Delta_k(N, \alpha) &= \sum_{n=0}^{N-1} e^{2\pi i q_k n \alpha}, \\ S(N, \alpha) &= \sum_{k \geq 1} r_k \operatorname{Re} \Delta_k(N, \alpha), \\ T(N, \alpha) &= \sum_{k \geq 1} r_k |\Delta_k(N, \alpha)|. \end{aligned} \quad (6)$$

Then

$$\begin{aligned}\Delta_k(N, \alpha) &= \frac{1 - e^{2\pi i q_k N \alpha}}{1 - e^{2\pi i q_k \alpha}}, \quad q_k \alpha \in \mathbb{R} \setminus \mathbb{Z}, \\ W(N, x, \alpha) &= \sum_{k \geq 1} r_k \operatorname{Re} \left[e^{2\pi i q_k x} \sum_{n=0}^{N-1} e^{2\pi i q_k n \alpha} \right] \\ &= \sum_{k \geq 1} r_k \operatorname{Re} \left[e^{2\pi i q_k x} \Delta_k(N, \alpha) \right],\end{aligned}$$

and the property (3) follows from the inequalities

$$\limsup_{N \rightarrow \infty} \frac{S(N, \alpha)}{\varphi(N)} > 0, \quad (7)$$

$$\limsup_{N \rightarrow \infty} \frac{T(N, \alpha)}{\varphi(N)} < \infty. \quad (8)$$

To get (7) and (8), we should first study the behavior of the sums $\Delta_k(N, \alpha)$. Essentially, if the fractional part of $q_k \alpha$ is of order $1/M$, then $\Delta_k(N, \alpha)$ behaves like N/M for N smaller than M , and is bounded by a constant times M for all N . After that, in an inductive process we approximate φ from below on an infinite sequence of points by a weighted sum of $\Delta_k(N, \alpha)$ with a lacunary sequence q_k and a suitable α .

Our first observation is as follows. Fix $n \geq 1$, suppose that the numbers $q_1, \dots, q_{n+1}$ satisfy condition (4), and define

$$k_n = \sum_{1 \leq s \leq n} \frac{q_n}{q_s} \in \mathbb{N}.$$

Claim. *Suppose that β belongs to the interval*

$$\mathcal{A}_n = \left\{ \beta : \frac{q_n}{q_{n+1}} \leq q_n \beta - k_n \leq \frac{2q_n}{q_{n+1}} \right\}. \quad (9)$$

Then

$$\left| \frac{\Delta_n(N, \beta)}{N} - 1 \right| \leq \frac{1}{2}, \quad 1 \leq N \leq \frac{q_{n+1}}{100q_n}, \quad (10)$$

$$|\Delta_n(N, \beta)| \leq \frac{q_{n+1}}{q_n}, \quad N \geq 1. \quad (11)$$

Proof. Applying the Taylor formula to the function $x \mapsto \exp ix$, and using that $\beta \in \mathcal{A}_n$ and $q_n/q_{n+1} \leq 1/100$, we obtain

$$\left| \frac{e^{2\pi i(q_n \beta - k_n)} - 1}{2\pi i(q_n \beta - k_n)} - 1 \right| \leq \left| \frac{(2\pi)^2 (q_n \beta - k_n)^2}{4\pi (q_n \beta - k_n)} \right| \leq \frac{1}{10}. \quad (12)$$

Furthermore, if $1 \leq N \leq q_{n+1}/(100q_n)$, then

$$\begin{aligned} \left| \frac{e^{2\pi i N(q_n \beta - k_n)} - 1}{2\pi i N(q_n \beta - k_n)} - 1 \right| &\leq \left| \frac{(2\pi)^2 N^2 (q_n \beta - k_n)^2}{4\pi N(q_n \beta - k_n)} \right| \\ &= \pi N(q_n \beta - k_n) \leq \frac{1}{10}. \end{aligned}$$

Hence,

$$\left| \frac{\Delta_n(N, \beta)}{N} - 1 \right| = \left| \frac{1 - e^{2\pi i N(q_n \beta - k_n)}}{N(1 - e^{2\pi i (q_n \beta - k_n)})} - 1 \right| \leq \frac{1}{2}, \quad 1 \leq N \leq \frac{q_{n+1}}{100q_n},$$

and (10) is proved.

Next,

$$|1 - e^{2\pi i N q_n \beta}| \leq 2, \quad N \geq 1.$$

Therefore,

$$|\Delta_n(N, \beta)| = \left| \frac{1 - e^{2\pi i N q_n \beta}}{1 - e^{2\pi i q_n \beta}} \right| \leq \frac{2}{|1 - e^{2\pi i (q_n \beta - k_n)}|}.$$

Using (12), we get

$$|\Delta_n(N, \beta)| \leq \frac{3}{2\pi(q_n \beta - k_n)} \leq \frac{q_{n+1}}{q_n}, \quad N \geq 1,$$

and (11) is proved. $\square$

Now, to obtain (7) and (8), we define q_k , r_k , and α in an inductive process. Without loss of generality we assume that $\varphi(1) \geq 2$. Set $S^0(N, \beta) = T^0(N, \beta) = 0$, $q_1 = 1$, $N_0 = M_0 = 1$, $\mathcal{A}_0 = (0, 1)$. On the induction step $p \geq 1$ we start with sequences $\{q_j\}_{1 \leq j \leq p}$, $\{r_j\}_{1 \leq j < p}$, $\{N_j\}_{0 \leq j < p}$, $\{M_j\}_{0 \leq j < p}$, and the interval $\mathcal{A}_{p-1}$ (defined by $\{q_j\}_{1 \leq j \leq p}$), such that for every $\beta \in \mathcal{A}_{p-1}$, the function S^{p-1} ,

$$S^{p-1}(N, \beta) = \sum_{1 \leq n < p} r_n \operatorname{Re} \Delta_n(N, \beta),$$

satisfies

$$S^{p-1}(N_j, \beta) \geq \frac{1}{100} \varphi(N_j) + 2^{-p}, \quad 1 \leq j < p, \quad (13)$$

and for every $\beta \in \mathcal{A}_{p-1}$ the function T^{p-1} ,

$$T^{p-1}(N, \beta) = \sum_{1 \leq n < p} r_n |\Delta_n(N, \beta)|,$$

satisfies

$$T^{p-1}(N, \beta) \leq 200 \varphi(N) - 2^{-p}, \quad N \geq 1, \quad (14)$$

$$T^{p-1}(N, \beta) \leq \frac{1}{100} \varphi(M_{p-1}), \quad N \geq 1. \quad (15)$$

By (6), for any β, N ,

$$|\Delta_p(N, \beta)| \leq N, \quad (16)$$

and we can choose r_p satisfying (5) such that for all $\beta \in \mathcal{A}_{p-1}$,

$$r_p(N + |\Delta_p(N, \beta)|) \leq 2^{-p-1}, \quad 1 \leq N \leq \max(M_{p-1}, N_{p-1}). \quad (17)$$

Since φ is increasing and $\lim_{N \rightarrow \infty} \varphi(N)/N = 0$, we can find

$$N_p > \max(M_{p-1}, N_{p-1})$$

such that

$$r_p N \leq \varphi(N), \quad N < N_p, \quad (18)$$

$$\varphi(N_p) \leq r_p N_p \leq \varphi(N_p) + 1. \quad (19)$$

Set $q_{p+1} = 100q_p N_p$. Then we define $\mathcal{A}_p \subset \mathcal{A}_{p-1}$ by the formula (9). In the estimates to follow we assume that $\beta \in \mathcal{A}_p$, and hence, by the Claim, (10) and (11) hold with $n = p$.

By (10) and (19),

$$r_p \operatorname{Re} \Delta_p(N_p, \beta) \geq \frac{1}{2} r_p N_p \geq \frac{1}{2} \varphi(N_p). \quad (20)$$

By (16) and (18),

$$r_p |\Delta_p(N, \beta)| \leq r_p N \leq \varphi(N), \quad N < N_p,$$

and by (11) and (19),

$$r_p |\Delta_p(N, \beta)| \leq 100 r_p N_p \leq 100(\varphi(N_p) + 1), \quad N \geq N_p. \quad (21)$$

Thus,

$$r_p |\Delta_p(N, \beta)| \leq 100(\varphi(N) + 1), \quad N \geq 1. \quad (22)$$

Now, by (13) and (17),

$$S^p(N_j, \beta) \geq \frac{1}{100} \varphi(N_j) + 2^{-p-1}, \quad 1 \leq j < p,$$

and by (15) and (20),

$$S^p(N_p, \beta) \geq \frac{1}{100} \varphi(N_p) + 2^{-p-1}.$$

Thus,

$$S^p(N_j, \beta) \geq \frac{1}{100} \varphi(N_j) + 2^{-p-1}, \quad 1 \leq j \leq p.$$

Furthermore, by (14) and (17),

$$T^p(N, \beta) \leq 200 \varphi(N) - 2^{-p-1}, \quad 1 \leq N \leq M_{p-1},$$

and by (15) and (22),

$$T^p(N, \beta) \leq 200 \varphi(N) - 1, \quad N > M_{p-1}.$$

Thus,

$$T^p(N, \beta) \leq 200 \varphi(N) - 2^{-p-1}, \quad N \geq 1.$$

Finally, by (15), (21), and by the condition that φ is unbounded, there exists M_p such that

$$T^p(N, \beta) \leq \frac{1}{100} \varphi(M_p), \quad N \geq 1.$$

This completes the induction step.

The intervals $\mathcal{A}_n$ constitute a nested family,

$$\bigcap_{n \geq 0} \mathcal{A}_n = \{\alpha\},$$

$$\alpha = \sum_{k \geq 1} \frac{1}{q_k},$$

and all the inequalities in the induction process are valid with $\beta = \alpha$.

We have

$$S(N, \alpha) = \lim_{p \rightarrow \infty} S^p(N, \alpha), \quad T(N, \alpha) = \lim_{p \rightarrow \infty} T^p(N, \alpha).$$

Then the properties (13) and (14) imply that

$$\frac{S(N_j, \alpha)}{\varphi(N_j)} \geq \frac{1}{100}, \quad j \geq 1,$$

$$\frac{T(N, \alpha)}{\varphi(N)} \leq 200, \quad N \geq 1,$$

and (7) and (8) follow. The theorem is proved. $\square$

The author is thankful to Leonid Polterovich and Misha Sodin for helpful remarks.

REFERENCES

- [1] G. D'Ambra, M. Gromov, *Lectures on transformation groups: geometry and dynamics*, in Surveys in Differential Geometry, supplement to J. Diff. Geom., 1 (1991), 19–112.
- [2] A. Borichev, *Distortion growth for iterations of diffeomorphisms of the interval*, manuscript, 2003.
- [3] A. Katok, B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Cambridge University Press, 1995.
- [4] L. Polterovich, *Slow symplectic maps, continued fractions and related stories*, In: Symplectic and Contact Topology: Interactions and Perspectives, eds. Y. Eliashberg, B. Khesin and F. Lalonde, Fields Institute Communications, **35** (2003).
- [5] L. Polterovich, M. Sodin, *A growth gap for diffeomorphisms of the interval*, preprint math.DS/0203066.

Alexander Borichev, Department of Mathematics,
University of Bordeaux I, 351, cours de la Libération, 33405 Talence, France